Parsing the electrophysiological signatures of spatially selective versus distributed attention in neuronal population data

BF


HOTCHKIS!
BRAIN INSTITUTION




Kiana Kazeminejad^{1,2}, Suprya Ghosh³, Jackson Cone^{1,2}

¹Departmnet of Biomedical Engineering, University of Calgary; ²Department of Psychology, University of Calgary; ³Department of Neurobiology, University of Chicago

Shifting attention from spatially selective to spatially distributed engages distinct neuronal population dynamics in visual area V4 via normalization circuits.






Lag (ms)

Lag (ms)

Changes in Pairwise Correlations Depend on Normalization Mechanisms



Future Directions

- Relate normalization model parameters to cluster identities in cross-correlation cluster analyses.
- Relate ISOMAP coordinate space to trial outcomes.

Implications

- Distributing attention to both visual hemifields engages neuronal mechanisms that differ considerably from established signatures of selective attention in visual area V4.
- Changes in neuronal correlations appear to depend on differential engagement of normalization circuitry.
- Prior work suggests that decorrelation of neuronal population activity plays an important role in the signatures of spatially selective attention that arise from attention. This work suggests that divergent population dynamics can yield equivalent levels of behavioural performance at the receptive field location of the recorded neurons.

